

Melanoma of the Skin Staging

7th EDITION

Definitions

Primary Tumor (T)

- TX** Primary tumor cannot be assessed (for example, curettaged or severely regressed melanoma)
- T0** No evidence of primary tumor
- Tis** Melanoma in situ
- T1** Melanomas 1.0 mm or less in thickness
- T2** Melanomas 1.01–2.0 mm
- T3** Melanomas 2.01–4.0 mm
- T4** Melanomas more than 4.0 mm

NOTE: a and b subcategories of T are assigned based on ulceration and number of mitoses per mm², as shown below:



T CLASSIFICATION	THICKNESS (mm)	ULCERATION STATUS/MITOSSES
T1	≤1.0	a: w/o ulceration and mitosis <1/mm ² b: with ulceration or mitoses ≥1/mm ²
T2	1.01–2.0	a: w/o ulceration b: with ulceration
T3	2.01–4.0	a: w/o ulceration b: with ulceration
T4	>4.0	a: w/o ulceration b: with ulceration

Regional Lymph Nodes (N)

- NX** Patients in whom the regional nodes cannot be assessed (for example, previously removed for another reason)
- N0** No regional metastases detected
- N1-3** Regional metastases based upon the number of metastatic nodes and presence or absence of intralymphatic metastases (in transit or satellite metastases)

NOTE: N1–3 and a–c subcategories assigned as shown below:

N CLASSIFICATION	NO. OF METASTATIC NODES	NODAL METASTATIC MASS
N1	1 node	a: micrometastasis ¹ b: macrometastasis ²
N2	2–3 nodes	a: micrometastasis ¹ b: macrometastasis ² c: in transit met(s)/satellite(s) without metastatic nodes
N3	4 or more metastatic nodes, or matted nodes, or in transit met(s)/satellite(s) with metastatic node(s)	



Distant Metastasis (M)

- M0** No detectable evidence of distant metastases
- M1a** Metastases to skin, subcutaneous, or distant lymph nodes
- M1b** Metastases to lung
- M1c** Metastases to all other visceral sites or distant metastases to any site combined with an elevated serum LDH

NOTE: Serum LDH is incorporated into the M category as shown below:

M CLASSIFICATION	SITE	SERUM LDH
M1a	Distant skin, subcutaneous, or nodal mets	Normal
M1b	Lung metastases	Normal
M1c	All other visceral metastases	Normal
	Any distant metastasis	Elevated

ANATOMIC STAGE/PROGNOSTIC GROUPS

Clinical Staging ³	Pathologic Staging ⁴					
	T	N	M			
Stage 0	Tis	N0	M0			
Stage IA	T1a	N0	M0			
Stage IB	T1b	N0	M0			
	T2a	N0	M0			
Stage IIA	T2b	N0	M0			
	T3a	N0	M0			
Stage IIB	T3b	N0	M0			
	T4a	N0	M0			
Stage IIC	T4b	N0	M0			
Stage III	Any T	≥ N1	M0			
			IIIA	T1-4a	N1a	M0
			IIIB	T1-4a	N2a	M0
		IIIC	T1-4b	N1a	M0	
		IIIC	T1-4b	N2a	M0	
		IIIC	T1-4a	N1b	M0	
	IIIC	T1-4a	N2b	M0		
	IIIC	T1-4a	N2c	M0		
	IIIC	T1-4b	N1b	M0		
	IIIC	T1-4b	N2b	M0		
	IIIC	T1-4b	N2c	M0		
	IIIC	Any T	N3	M0		
Stage IV	Any T	Any N	M1			

Notes

- ¹ Micrometastases are diagnosed after sentinel lymph node biopsy and completion lymphadenectomy (if performed).
- ² Macrometastases are defined as clinically detectable nodal metastases confirmed by therapeutic lymphadenectomy or when nodal metastasis exhibits gross extracapsular extension.
- ³ Clinical staging includes microstaging of the primary melanoma and clinical/radiologic evaluation for metastases. By convention, it should be used after complete excision of the primary melanoma with clinical assessment for regional and distant metastases.
- ⁴ Pathologic staging includes microstaging of the primary melanoma and pathologic information about the regional lymph nodes after partial or complete lymphadenectomy. Pathologic Stage 0 or Stage IA patients are the exception; they do not require pathologic evaluation of their lymph nodes.



Financial support for AJCC 7th Edition Staging Posters provided by the American Cancer Society